

# Challenges to Accessing Reproductive Health Services in New Jersey

Dulce Gonzalez and Emily M. Johnston


New data from the 2024–25 [Reproductive Health Experiences and Access \(RHEA\) survey](#) offer insight into access to reproductive health services [in New Jersey](#). Reproductive health care includes many services, such as pelvic exams and screening for cervical cancer, care for irregular or painful periods, birth control, fertility assistance, gender-affirming care, and care for menopause symptoms. Factors like cost, insurance barriers, and difficulty securing an appointment can prevent people from accessing the care they want and need. New Jersey has a “[very protective](#)” abortion policy environment, according to the Guttmacher Institute, and [has expanded Medicaid eligibility](#) to childless adults. Our analyses include responses from 2,801 [women<sup>1</sup> ages 18 to 49](#) residing in New Jersey.

## NEARLY 1 IN 10 RESPONDENTS DELAYED OR HAD TROUBLE GETTING THEIR DESIRED BIRTH CONTROL

In 2024–25, 8 percent of respondents reported they delayed or had trouble getting the birth control method they wanted in the past year. Compared with the US overall (analysis of the RHEA survey, data not shown), New Jersey respondents were slightly more likely to report delay or trouble getting birth control (8 percent versus 7 percent). Among the demographic groups we examined (figure 1), New Jersey respondents ages 45 to 49 (3 percent) were less likely than those ages 18 to 24 (11 percent) and those ages 25 to 44 (9 percent) to report delay or trouble getting birth control. People with a disability (17 percent), less than a bachelor’s degree (11 percent), public or no health insurance (12 percent), or who identify as LGBTQIA+ (18 percent) also reported higher rates than others.<sup>2</sup>

FIGURE 1

### The Share Reporting Delay or Trouble Getting Birth Control in the past 12 Months Varies by Group



URBAN INSTITUTE

Source: 2024–25 Reproductive Health Experiences and Access survey.

Notes: Samples for each subgroup range from 303 to 2,426 people assigned female at birth, ages 18 to 49, residing in New Jersey. Demographic groups examined include age, disability status, educational attainment, insurance status, income, sexual orientation and gender identity, race/ethnicity, rurality, and nativity. Only groups for which the sample size is at least 200, the difference is at least 5 percentage points, and the difference is statistically significant at  $p < 0.05$  are presented.

<sup>1</sup> The RHEA survey population consists of people assigned female at birth.


<sup>2</sup> All differences reported between New Jersey and the US and across subgroups are significant at the  $p < 0.05$  level.

## AROUND 3 IN 10 NEEDED BUT DID NOT GET ONE OR MORE TYPES OF REPRODUCTIVE HEALTH CARE SERVICES

Overall, 29 percent of respondents needed but did not get one or more types of care in the last 12 months. Figure 2 shows that about 16 percent of respondents needed but did not get preventive gynecologic care, and 9 percent did not get needed care for irregular or painful periods. Care for premenstrual dysphoric disorder, gender-affirming care, and care for loss of sexual desire or dysfunction were reported as unneeded by 90 percent or more of respondents. Among those who did need these types of care, at least half did not receive it.

FIGURE 2

### Sixteen Percent of New Jersey Respondents Did Not Get Needed Preventive Gynecologic Care



URBAN INSTITUTE

Source: 2024–25 Reproductive Health Experiences and Access survey.

Notes: Samples for each type of care range from 2,688 to 2,751 people assigned female at birth, ages 18 to 49, residing in New Jersey. For each type of care, estimates exclude respondents with missing responses and those who responded “don’t know” or “prefer not to answer.” Percentages may not sum to 100 due to rounding.

Compared with the US overall (data not shown), New Jersey respondents were slightly less likely to report needing but not getting one or more types of reproductive health care (29 percent versus 30 percent). Among demographic groups we examined (data not shown), those ages 25 to 44 (31 percent) and 45 to 49 (31 percent) were more likely than those ages 18 to 24 (21 percent) to report not getting one or more needed reproductive health services. People with disabilities (42 percent), less than a bachelor’s degree (32 percent), public or no health insurance (36 percent), or who identify as LGBTQIA+ (37 percent) were also more likely than others to report not getting one or more needed reproductive health services.

#### ABOUT THE RHEA STUDY

This fact sheet is part of the [RHEA Study](#), a multiyear assessment of reproductive health access, experiences, and preferences following the 2022 US Supreme Court decision in *Dobbs v. Jackson Women’s Health Organization*, especially among people facing barriers to care. For this study, the Urban Institute, the Reproductive Equity Action Lab at the University of Wisconsin–Madison, and SisterSong Women of Color Reproductive Justice Collective are collaborating to produce evidence-based research findings through a large national survey and qualitative data collection in 13 states.